



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

and center at the origin. Since each double point counts as two points the circle cuts the septic curve in 20 points. But this is impossible, hence there is no double point except at the origin. Therefore  $\delta=5$ . Then from Plücker's equations:  $m=32$ ,  $\iota=75$ ,  $\tau=380$ . Hence the curve

$$5x^4y - 10x^2y^3 + y^5 + c(x^2 + y^2)(x^5 - 10x^3y + 5xy^4) = 0, \quad c \neq 0.$$

is of class 32, is non-cuspidal, and has five double points at the origin, 75 points of inflection of which five are at the origin, and 380 double tangents. It is obvious that not all the singularities are real.

---

## DEPARTMENTS.

---

### SOLUTIONS OF PROBLEMS.

---

#### ALGEBRA.

---

330. Proposed by R. D. CARMICHAEL, Princeton, N. J.

An important function in the Theory of Numbers is one defined thus:  $f(x)=1$  when  $x>0$ ,  $f(x)=0$  when  $x=0$ ,  $f(x)=-1$  when  $x<0$ . Two analytic expressions for  $f(x)$  are the following:

$$f(x) = \lim_{n \rightarrow \infty} x^{1/(2n-1)}, \quad n=1, 2, \dots; \quad f(x) = \lim_{n \rightarrow \infty} \frac{(x+1)^n - (x+1)^{-n}}{(x+1)^n + (x+1)^{-n}}, \quad x > -1.$$

It is required to find other non-trigonometric analytic expressions for this function. (There are several representations of  $f(x)$  by means of trigonometric functions.)

No solution of this problem has been received.

331. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Extract the square root of  $21+6\sqrt{2}+2\sqrt{21}-6\sqrt{3}-6\sqrt{7}-2\sqrt{6}-2\sqrt{14}$  and also of  $4\sqrt{2}+2\sqrt{6}-9-4\sqrt{3}$ .

Solution by S. G. BARTON, Ph. D., Clarkson School of Technology, Potsdam, N. Y., and J. SCHEFFER, A.M., Hagerstown, Md.

(a) Assume the root to be of the form

$$a\sqrt{2}+b\sqrt{3}+c\sqrt{7}+d.$$

Squaring and comparing coefficients, we have

$$ab=-1, ac=-1, ad=3, bc=1, bd=-3, cd=-3, \\ 2a^2+3b^2+7c^2+d^2=21.$$

Whence  $a=-1, b=1, c=1, d=-3$ , and the root is  $\sqrt[3]{3-\sqrt{2}+\sqrt{7}-3}$ .

For second expression, (b), assume root of form  $\sqrt[3]{-1(a+b\sqrt{3}+c\sqrt{2})}$ . Squaring and comparing coefficients, we have,

$$ab=2, ac=-2, bc=-1, a^2+3b^2+2c^2=9.$$

Whence  $a=2, b=1, c=-1$ , and the root is  $\sqrt[3]{-1(2+\sqrt{3}-\sqrt{2})}$ .

Solved similarly by V. M. Spunar and Levi S. Shively.

## II. Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

The method used here was suggested to me by Mr. Githens in a solution for cube root. It is considerably simpler than methods ordinarily used.

$$(a) \text{ Let } A=[21+6\sqrt{2}+2\sqrt{21}-(6\sqrt{3}+6\sqrt{7}+2\sqrt{6}+2\sqrt{14})]^{\frac{1}{3}} \\ =\sqrt[3]{(m-n)}.$$

$$\text{Then } \sqrt[3]{(m-n)(m+n)}=[157+12\sqrt{2}-(4\sqrt{21}+24\sqrt{42})]^{\frac{1}{3}}=\sqrt[3]{(p-q)}.$$

$$\sqrt[3]{(p-q)(p+q)}=(409-264\sqrt{2})^{\frac{1}{3}}=\sqrt[3]{(r-s)}.$$

$$\sqrt[3]{(r-s)(r+s)}=167.$$

Let  $\sqrt[3]{(r-s)}=\sqrt[3]{(x+167)}-\sqrt[3]{x}$ . Squaring both members and equating rational terms,  $2x+167=409$ , or  $x=121$ .

$$\therefore \sqrt[3]{(r-s)}=12\sqrt[3]{2}-11.$$

$$\text{Let } \sqrt[3]{(p-q)}=\sqrt[3]{(x+12\sqrt[3]{2}-11)}-\sqrt[3]{x}.$$

$$\text{Then } 2x+12\sqrt[3]{2}-11=157+12\sqrt[3]{2}, \text{ or } x=84.$$

$$\therefore \sqrt[3]{(p-q)}=\sqrt[3]{(73+12\sqrt[3]{2})}-2\sqrt[3]{21}.$$

$$\sqrt[3]{[(73+12\sqrt[3]{2})(73-12\sqrt[3]{2})]}=71.$$

$$\text{Hence } \sqrt[3]{(73+12\sqrt[3]{2})}=\sqrt[3]{(x+71)}+\sqrt[3]{x}.$$

$$\text{From this, } x=1 \text{ and } \sqrt[3]{(73+12\sqrt[3]{2})}=6\sqrt[3]{2}+1.$$

$$\sqrt[3]{(p-q)}=6\sqrt[3]{2}+1-2\sqrt[3]{21}.$$

$$\text{Let } \sqrt[3]{(m-n)}=\sqrt[3]{(x+6\sqrt[3]{2}+1-2\sqrt[3]{21})}-\sqrt[3]{x}.$$

$$\text{Then } 2x+6\sqrt[3]{2}+1-2\sqrt[3]{21}=21+6\sqrt[3]{2}+2\sqrt[3]{21}.$$

$$\therefore x=10+2\sqrt[3]{21}. \quad \therefore \sqrt[3]{(m-n)}=\sqrt[3]{(11+6\sqrt[3]{2})}-\sqrt[3]{(10+2\sqrt[3]{21})}.$$

$$\sqrt[3]{[(11+6\sqrt[3]{2})(11-6\sqrt[3]{2})]}=7, \text{ and } \sqrt[3]{(11+6\sqrt[3]{2})}=\sqrt[3]{(x+7)}+\sqrt[3]{x} \dots (1).$$

$$\sqrt[3]{[(10+2\sqrt[3]{21})(10-2\sqrt[3]{21})]}=4, \text{ and } \sqrt[3]{(10+2\sqrt[3]{21})}=\sqrt[3]{(x+4)}+\sqrt[3]{x} \dots (2).$$

From (1),  $x=2$ ; from (2),  $x=3$ .

$$\therefore \sqrt[3]{(11+6\sqrt[3]{2})}=3+\sqrt[3]{2}, \quad \sqrt[3]{(10+2\sqrt[3]{21})}=\sqrt[3]{7}+\sqrt[3]{3}.$$

$$\therefore \sqrt[3]{(m-n)}=3+\sqrt[3]{2}-\sqrt[3]{7}-\sqrt[3]{3}.$$

(b) Let  $(-9-4\sqrt{3}+4\sqrt{2}+2\sqrt{6})^{\frac{1}{2}} = \sqrt{-m+n}$ .

$$\sqrt{[(-m+n)(-m-n)]} = \sqrt{[(-m)^2 - n^2]} = \sqrt{[73+40\sqrt{3}]} = \sqrt{[p+q]}.$$

$$\sqrt{[(p+q)(p-q)]} = 23.$$

Let  $\sqrt{p+q} = \sqrt{x+23} + \sqrt{x}$ . Then  $2x+23=73$ , or  $x=25$ .

$$\therefore \sqrt{p+q} = 4\sqrt{3}+5.$$

Let  $\sqrt{n-m} = -\sqrt{x+4\sqrt{3}+5} + \sqrt{x}$ . Then  $2x+4\sqrt{3}+5 = -9-4\sqrt{3}$ , or  $x = -7-4\sqrt{3}$ .

$$\therefore \sqrt{n-m} = \sqrt{-7-4\sqrt{3}} - \sqrt{-2}.$$

$$\sqrt{[(-7-4\sqrt{3})(-7+4\sqrt{3})]} = 1.$$

Let  $\sqrt{-7-4\sqrt{3}} = \sqrt{x+1} + \sqrt{x}$ ; then  $2x+1 = -7$ ,  $x = -4$ .

$$\therefore \sqrt{-7-4\sqrt{3}} = \sqrt{-3} + 2\sqrt{-1}.$$

$$\therefore \sqrt{n-m} = \sqrt{-3} + 2\sqrt{-1} - \sqrt{-2}.$$

Also solved by G. I. Hopkins and A. H. Holmes.

332. Proposed by C. N. SCHMALL, New York City.

Solve the quadratic,  $x^2 + ax + b = 0$ , without completing the square.

Solution by ARTEMAS MARTIN, LL. D., Washington, D. C.

Assume  $y - \frac{1}{2}a = x$ , and substitute in the given quadratic and it becomes

$$(y - \frac{1}{2}a)^2 + a(y - \frac{1}{2}a) + b = 0, \text{ or } y^2 - \frac{1}{4}a^2 + b = 0;$$

whence  $y = \pm \sqrt{(\frac{1}{4}a^2 - b)}$  and  $x = \pm \sqrt{\frac{1}{4}a^2 - b} - \frac{1}{2}a$ .

See *Mathematical Magazine*, Vol. I, No. 9 (January, 1884), p. 146.

Solved similarly by V. M. Spunar, Levi S. Shively and the Proposer.

Professor Hopkins, in his solution, made use of the principle that the sum of the roots is equal to the coefficient of  $x$  with sign changed, and the product of the roots is equal to the final term.

S. Lefschetz sent in solutions of 327 and 328 too late for credit in last issue.


## GEOMETRY.

356. Proposed by G. I. HOPKINS, Manchester, N. H.

Required to construct a triangle having given, base, vertical angle, and difference of other two sides.

I. Solution by J. M. ARNOLD, Crompton, R. I.

Let  $x$  = the longer side, then  $x-d$  = the shorter side. Let  $A$  = the vertical angle. Then



$$x^2 - (x \cos A)^2 = (x-d)^2 - (b-x \cos A)^2$$

which gives

$$(2b \cos A - 2d)x = b^2 - d^2.$$

Dividing by 4 and putting in the form of a proportion